

MURI_MOXA_seashr

** PROFILE FOR NO2 IS BASED ON THE 98th percentile by Seasonal -Hourly.
 ** DATA COLLECTED AT THE MURI -MOXA SITE, YEARS 2009-2012.
 **
 **
 ** BACKGROUND CONCENTRATIONS ARE IN PPB
 SO BACKUNIT PPB
 ** Winter (December, January, and February):
 SO BACKGRND SEASHR 10.2 8.7 10.1 7.3 9.8 8.9 11.4 13.8 9 7.9 8.7 7 6.1 5.9 7.1 8.7
 7.2 9.1 9.9 9 10.4 11.1 11 10.6
 ** Spring (March, April, and May):
 SO BACKGRND SEASHR 8.3 6.8 7.7 7.4 9 8.7 7.2 6.7 6.4 5.3 3.5 2.6 2.9 3 2.3 2.9 3 2.5
 3.2 3.3 5 5.3 7 8.9
 ** Summer (June, July, and August):
 SO BACKGRND SEASHR 6.7 6.8 6.6 7.6 7.9 6.7 6.7 6.6 6.3 7.3 5.5 3.6 2.8 2.3 2.1 2 2.2
 2.5 3.3 3.2 4 5.4 6.5 7.7
 ** Fall (September, October, and November):
 SO BACKGRND SEASHR 8.8 11.3 9.3 9.5 9.2 10.5 10.1 9.2 7.7 7 8.8 5.3 5.8 6.5 4.3 5.8
 6 5.4 5.6 5.7 7.5 9.3 9.7 9.4